

Synthesis and characterization of the pentamolybdate ion, $\text{Mo}_5\text{O}_{17}\text{H}_3^-$

M. Filowitz, W. G. Klemperer, and W. Shum

J. Am. Chem. Soc., **1978**, 100 (8), 2580-2581 • DOI: 10.1021/ja00476a069

Downloaded from <http://pubs.acs.org> on December 5, 2008

More About This Article


The permalink <http://dx.doi.org/10.1021/ja00476a069> provides access to:

- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article



ACS Publications
High quality. High impact.

certed mechanism can compete with radical cleavage except possibly for the worst leaving groups.



Relative rates of β cleavage can be obtained directly from the product ratios, given the plausible assumption that k_s is largely independent of X.¹⁶ Recent measurements¹⁷ indicate that $k_s = 1 \times 10^7 \text{ s}^{-1}$; so a k_{-X} value of $2.7 \times 10^5 \text{ s}^{-1}$ is indicated for X = *n*-BuS. Actual rate constants for β elimination in solution have not been reported previously. The 7.5-kcal activation energy estimated for β -bromoethyl radical¹⁸ is certainly consistent with our estimated $k_{-Br} = 1 \times 10^8 \text{ s}^{-1}$, given an *A* factor of 10^{13} s^{-1} .

Comparison of the relative k_{-X} values is revealing. As expected, $I > Br > Cl$ and $PhS \gg n\text{-BuS}$ ¹⁹ as leaving groups. The order $RSO \gg RS$, RSO_2 confirms Kice's suggestion²⁰ that sulfinyl radicals have the greatest relative kinetic stability of the three. That $SCN \gg SCOCH_3$ may reflect the known weak conjugative stabilization in α -keto radicals²¹ and further indicates that $\delta\text{-X}$ leaves as a radical rather than an anion.

The most surprising result is that Cl is eliminated more rapidly than thyl radicals. Kineticists usually assume a much smaller value of k_{-X} for β -chloroethyl²² than we estimate. Therefore it is possible that the cyclic concerted mechanism contributes for X = Cl. However, the amount of Cl loss in the radical addition of mercaptans to allyl chloride²³ is consistent with our separate observations that loss of $\delta\text{-Cl}$ and trapping by mercaptans^{14a} compete comparably with type II cleavage. It is not inconceivable that studies on monoradicals have indicated too low a value for k_{-Cl} because of rapid reverse addition. Our k_{-X} values are *minimum* values uncorrected for reverse addition. However, since the diradicals can generate caged radical pairs, in-cage radical-radical reactions are probably faster even than addition of Cl atoms to double bonds and thus minimize the effects of reversibility on relative k_{-X} values.

References and Notes

- (1) We thank the National Science Foundation for partial support of this work.
- (2) P. J. Wagner, *Acc. Chem. Res.*, **4**, 168 (1971).
- (3) (a) For a review, see P. S. Skell and K. J. Shea in "Free Radicals", J. K. Kochi, Ed., Wiley-Interscience, New York, N.Y., Vol. II, p. 809; (b) D. J. Edge and J. K. Kochi, *J. Am. Chem. Soc.*, **94**, 6485 (1972); (c) K. S. Chen, P. J. Krusic, P. Meakin, and J. K. Kochi, *J. Phys. Chem.*, **78**, 2014 (1974); (d) R. V. Lloyd and D. E. Wood, *J. Am. Chem. Soc.*, **97**, 5986 (1975); (e) D. Griller and K. U. Ingold, *ibid.*, **96**, 6715 (1974).
- (4) P. J. Wagner and A. E. Kempainen, *J. Am. Chem. Soc.*, **94**, 7495 (1972).
- (5) III was collected by preparative GC and identified by comparison of its NMR and mass spectra with those of independently synthesized material. None of the ketones V-X produce III in the dark or during analysis.
- (6) NMR spectra indicated no *N*-alkylpyridinium halide formation in our solution. Pyridine does not affect triplet state decay: P. J. Wagner, T. Jellinek, and A. E. Kempainen, *J. Am. Chem. Soc.*, **94**, 7512 (1972).
- (7) P. J. Wagner, I. E. Kochevar, and A. E. Kempainen, *J. Am. Chem. Soc.*, **94**, 7489 (1972).
- (8) (a) C. Walling, "Free Radicals in Solution", Wiley, New York, N.Y., 1957, p. 302; (b) E. S. Huyser, "Free-Radical Chain Reactions", Wiley-Interscience, New York, N.Y., 1970, p. 212.
- (9) P. J. Kropp, G. S. Poindexter, N. J. Pienta, and D. C. Hamilton, *J. Am. Chem. Soc.*, **98**, 8135 (1976).
- (10) C. Walling and W. Helmreich, *J. Am. Chem. Soc.*, **81**, 1144 (1959).
- (11) J. B. Gutenplan and S. G. Cohen, *J. Am. Chem. Soc.*, **95**, 200 (1973).
- (12) (a) R. M. Topping and N. Kharasch, *J. Org. Chem.*, **27**, 4353 (1962); (b) D. Barnard and E. J. Percy, *Chem. Ind. (London)*, 1332 (1960).
- (13) (a) A. Padwa, *J. Am. Chem. Soc.*, **87**, 4205 (1965); (b) N. Shimizu, M. Ishikawa, K. Ishikura, and S. Nishida, *ibid.*, **96**, 6456 (1974); (c) P. J. Wagner and K.-C. Liu, *ibid.*, **96**, 5952 (1974).
- (14) (a) P. J. Wagner and R. G. Zepp, *J. Am. Chem. Soc.*, **94**, 287 (1972); (b) J. Grotewold, C. M. Previtali, D. Soria, and J. C. Scaiano, *J. Chem. Soc., Chem. Commun.*, 207 (1973); (c) M. Hamity and J. C. Scaiano, *J. Photochem.*, **4**, 229 (1975); (d) H. E. O'Neal, R. G. Miller, and E. Gunderson, *J. Am. Chem. Soc.*, **96**, 3351 (1974); (e) J. C. Scaiano, *ibid.*, **99**, 1494 (1977); (f) R. D. Small and J. C. Scaiano, *J. Phys. Chem.*, **81**, 828 (1977).
- (15) P. J. Wagner and R. G. Zepp, *J. Am. Chem. Soc.*, **93**, 4958 (1971).
- (16) If the rate-determining step for triplet generated diradical reaction is $T \rightarrow S$ intersystem crossing,^{14a} heavy-atom effects of δ substituents might in-

crease k_s .

- (17) R. D. Small, Jr., and J. C. Scaiano, *Chem. Phys. Lett.*, **50**, 431 (1977).
- (18) (a) R. Barker and A. Maccoll, *J. Chem. Soc.*, 2839 (1963); (b) P. I. Abell and R. S. Anderson, *Tetrahedron Lett.*, 3727 (1964).
- (19) D. N. Hall, A. A. Oswald, and K. Griesbaum, *J. Org. Chem.*, **30**, 3829 (1965).
- (20) J. L. Kice in ref 3a, pp 715-718.
- (21) (a) G. A. Russell and J. Lokensgard, *J. Am. Chem. Soc.*, **89**, 5059 (1967); (b) K. D. King, D. M. Golden, and S. W. Benson, *ibid.*, **92**, 5541 (1970).
- (22) (a) R. Eckling, P. Goldfinger, G. Huybrechts, G. Martens, L. Meyers, and S. Smoes, *Chem. Ber.*, **93**, 3014 (1960); (b) P. B. Ayscough, A. J. Cocker, F. S. Dainton, S. Hirst, and M. Weston, *Proc. Chem. Soc. London*, 244 (1961).
- (23) D. N. Hall, *J. Org. Chem.*, **32**, 2082 (1967).

Peter J. Wagner,* James H. Sedon
Michael J. Lindstrom


Department of Chemistry, Michigan State University
East Lansing, Michigan 48824

Received May 23, 1977

Synthesis and Characterization of the Pentamolybdate Ion, $Mo_5O_{17}H^{3-}$

Sir:

We recently reported the synthesis and characterization of the dimolybdate ion, $Mo_2O_7^{2-}$, as a tetrabutylammonium salt.¹ This compound contains the most basic polymolybdate ion isolated to date and is reactive toward a wide variety of reagents. We report here its reaction with water to form a labile pentamolybdate ion:



The instability of this novel, protonated isopolymolybdate in nonaqueous solution provides new insights into the kinetics and mechanism of polymolybdate transformations.

Although addition of water to a CH_3CN solution of $[(n\text{-}C_4H_9)_4N]_2Mo_2O_7$ (**1**) produces no significant change in its solution IR spectrum,¹ the pentamolybdate ion can be synthesized according to eq 1 by forcing the equilibrium to the right using two different approaches. In the first approach, the low solubility of the pentamolybdate ion in $CH_3CN/(C_2H_5)_2O$ is exploited. Slow addition ($\sim 15 \text{ mL/min}$) of 80 mL of $(C_2H_5)_2O$ to a clear solution of 1.0 g of **1** and 1.0 mL of H_2O in 10 mL of CH_3CN with rapid stirring yields 190 mg of an amorphous precipitate having the empirical formula² $[(n\text{-}C_4H_9)_4N]_3Mo_5O_{17}H$ (**2**). In the second approach, the high solubility of $[(n\text{-}C_4H_9)_4N]_2MoO_4$ in H_2O relative to **1** and **2** is exploited. When a suspension of 1.0 g of **1** in 10 mL of pH 5-6 aqueous HCl is stirred for $\sim 1 \text{ min}$ and filtered, compound **2** (460 mg) is obtained as an amorphous powder.

Structural characterization of **2** is rendered difficult by its instability in solution (see below) which has thus far ruled out recrystallization, conductivity measurements, and the measurement of ^{17}O NMR parameters. Nonetheless, elemental analysis² and IR spectroscopy strongly suggest the anion structure for **2** shown in Figure 1, where MoO_4^{2-} and OH^- groups are connected to opposite sides of an Mo_4O_{12} ring by weak ($>2.2 \text{ \AA}$) Mo-O bonds yielding the structural formula³ $(Mo_4O_{12})(OH)(MoO_4)$. This structure is closely related to the $(CH_3)_2AsMo_4O_{15}H^{2-}$ structure⁴ by replacement of the bidentate MoO_4^{2-} subunit with a bidentate $(CH_3)_2AsO_2^-$ group.⁵ Structural isomorphism between the anions in **2** and $[(n\text{-}C_4H_9)_4N]_2(CH_3)_2AsMo_4O_{15}H$ (**3**) should imply similar IR spectra for the two compounds in the 500-4000- cm^{-1} region with the exception of absorptions characteristic of the MoO_4^{2-} and $(CH_3)_2AsO_2^-$ subunits. Such a similarity is in fact observed. Both **2** and **3** exhibit sharp OH absorptions at 3610 cm^{-1} and their spectra in the 500-1000- cm^{-1} region (see Figure 2a and 2b) bear a striking resemblance outside of the 725-850- cm^{-1} region. In the 725-850- cm^{-1} region, **3** displays





Figure 1. Proposed, C_{2v} idealized structure for $\text{Mo}_5\text{O}_{17}\text{H}^{3-}$. Small filled circles represent molybdenum atoms, large open circles represent oxygen atoms, and the shaded large circle represents a hydroxyl group. Molybdenum-oxygen distances $< 1.75 \text{ \AA}$ are represented by double lines, $1.75 \text{ \AA} < d_{\text{Mo-O}} < 1.95 \text{ \AA}$ by single lines, and $d_{\text{Mo-O}} > 2.2 \text{ \AA}$ by dotted lines.

characteristic⁸ As-O absorptions at 803 and 826 cm^{-1} and **2** displays absorptions at 786 and 740 cm^{-1} which may be assigned to tetrahedral molybdate.

When **2** is dissolved in CH_3CN , its solution IR spectrum in the $700\text{--}1000\text{-cm}^{-1}$ region (see Figure 2c) taken < 2 min after dissolution shows absorptions characteristic of **1**¹ at 880 and 786 cm^{-1} and $\alpha\text{-}[(n\text{-C}_4\text{H}_9)_4\text{N}_4\text{Mo}_8\text{O}_{26}]$ (**4**)^{9,10} at 950 , 920 , 905 , 850 , 808 , and 740 cm^{-1} , but no pattern of absorptions characteristic of **2**. An analogous spectrum in $1,2\text{-C}_2\text{H}_4\text{Cl}_2$ (see Figure 2d), however, shows absorptions due to pentamolybdate **2** at 954 , 934 , 913 , 786 , and 740 cm^{-1} in addition to weaker absorptions characteristic of di- and octamolybdate. When this dichloroethane solution is allowed to equilibrate, precipitation of **4** is accompanied by an increase in intensity of the absorptions for **1** and **4** and a concomitant decrease in intensity of the absorptions for **2** (see Figure 2e). These observations imply the following equilibrium



which is rapid and favors decomposition products in CH_3CN , but is slower and less displaced toward decomposition products in $1,2\text{-C}_2\text{H}_4\text{Cl}_2$.

Two features of the $(\text{MoO}_4^{2-})(\text{OH}^-)(\text{Mo}_4\text{O}_{12})$ structure account for its instability in solution. First, the presence of an OH^- subunit offers the potential for generation of protons in solution capable of reacting with basic molybdate species to form condensation products. ^1H NMR measurements confirm the lability of this proton in that addition of CH_3OH to a solution of **2** in $1,2\text{-C}_2\text{H}_4\text{Cl}_2$ causes the δ 2.12 ppm resonance for **2** to disappear.¹² Second, the presence of a weakly bound MoO_4^{2-} subunit implies facile dissociation of the MoO_4^{2-} group in a fashion analogous to that observed for $(\text{MoO}_4^{2-})_2(\text{Mo}_6\text{O}_{18})$.³ Dynamic ^{17}O NMR experiments designed to confirm this lability have been unsuccessful since room temperature spectra cannot be accumulated before equilibration is complete and low temperature spectra yield resonances broadened beyond recognition by quadrupole relaxation. These two features suggest a simple, four-step mechanism for the decomposition reaction given in eq 2:






Figure 2. IR spectra of (a) $[(n\text{-C}_4\text{H}_9)_4\text{N}]_3\text{Mo}_5\text{O}_{17}\text{H}$ (**2**) in KBr pellet, (b) $[(n\text{-C}_4\text{H}_9)_4\text{N}]_2(\text{CH}_3)_2\text{AsMo}_4\text{O}_{15}\text{H}$ (**3**) in KBr pellet, (c) **2** in CH_3CN at equilibrium, (d) **2** in $1,2\text{-C}_2\text{H}_4\text{Cl}_2$ < 2 min after dissolution (e) **2** in $1,2\text{-C}_2\text{H}_4\text{Cl}_2$ 30 min after dissolution. Spectra c-e were measured in ZnS (IRTRAN-2) cells.



In the initial step, dissociation and deprotonation yield monomolybdate ions, protons, and $(\text{O}^{2-})(\text{Mo}_4\text{O}_{12})$ fragments. The protons are utilized in the second step to condense monomolybdate ions into dimolybdate ions. In the third step, the $(\text{O}^{2-})(\text{Mo}_4\text{O}_{12})$ fragments dimerize to form $[(\text{O}^{2-})(\text{Mo}_4\text{O}_{12})]_2$, i.e., $\beta\text{-Mo}_8\text{O}_{26}^{4-}$. Finally, $\beta\text{-Mo}_8\text{O}_{26}^{4-}$ undergoes facile isomerization to $\alpha\text{-Mo}_8\text{O}_{26}^{4-}$ via the mechanism proposed in ref 9.

Acknowledgments. We are grateful to the National Science Foundation for support of this research.

References and Notes

- V. W. Day, M. F. Fredrich, W. G. Klemperer, and W. Shum, *J. Am. Chem. Soc.*, **99**, 6146 (1977).
- Calculated (found) weight percentages: C, 38.95 (39.02); H, 7.42 (7.50); N, 2.84 (2.91); Mo, 32.41 (32.30).
- V. W. Day, M. F. Fredrich, W. G. Klemperer, and W. Shum, *J. Am. Chem. Soc.*, **99**, 952 (1977).
- K. M. Barkigia, L. M. Rajković, M. T. Pope, and C. O. Quicksall, *J. Am. Chem. Soc.*, **97**, 4146 (1975).
- A similar relationship is known to exist between the $\alpha\text{-Mo}_8\text{O}_{26}^{4-}$ and $(\text{CH}_3)_2\text{As}_2\text{Mo}_8\text{O}_{24}^{4-}$ structures^{3,6,7} which are related by replacement of a tridentate MoO_4^{2-} subunit by a tridentate $\text{CH}_3\text{AsO}_3^{2-}$ unit.
- J. Fuchs and H. Hartl, *Angew. Chem., Int. Ed. Engl.*, **15**, 375 (1976).
- W. Kwak, L. M. Rajković, J. K. Stalick, M. T. Pope, and C. O. Quicksall, *Inorg. Chem.*, **15**, 2778 (1976).
- A. Simon and H. D. Schumann, *Z. Anorg. Allg. Chem.*, **399**, 97 (1973); H. V. Gröndler, H. D. Schumann, and E. Steger, *J. Mol. Struct.*, **21**, 149 (1974).
- W. G. Klemperer and W. Shum, *J. Am. Chem. Soc.*, **98**, 8291 (1976).
- The solid state IR and Raman spectra of $[(n\text{-C}_4\text{H}_9)_4\text{N}]_4\text{Mo}_8\text{O}_{26}$ given in ref 11 are not characteristic of pure **3** but are characteristic of hydrated **3** obtained from either unrecrystallized aqueous precipitates or pure **3** which has been exposed to atmospheric water for 1 to 3 days.
- J. Fuchs and I. Brudgam, *Z. Naturforsch. B*, **32**, 853 (1977).
- The ^1H NMR spectrum of 70 mg of **2** in 0.6 mL of $1,2\text{-C}_2\text{H}_4\text{Cl}_2$ at 30°C displays a δ 2.12 ppm resonance both in freshly prepared and equilibrated solutions. This resonance disappears upon addition of a drop of CH_3OH .
- Fellow of the Alfred P. Sloan Foundation, 1976–1978.

M. Filowitz, W. G. Klemperer,^{*13} W. Shum

Department of Chemistry, Columbia University
New York, New York 10027

Received November 21, 1977