




SCIENTIFIC TOPICS

Fundamentals for emergent systems

Recent results published independently by the IRN-POM consortium revealed that solution-chemistry of POMs was not fully understood and that the fine control of the non-covalent interactions open new perspectives in the manipulation of POMs at the supramolecular level.

These systems have a structural organization defined by certain forces (hydrogen bonding, π -stacking, electrostatic, chaotropic, hydrophobic, metal coordination...), that could respond to




external and internal triggers such as light, redox, temperature, or pH. Joint expertise is therefore a prerequisite for the bottom-up construction, whether directed or not, of hierarchical structures or molecular devices that will bring out emergent properties or enable the synergistic combination of properties.

Advanced analytical tools, data treatment and theoretical modelling

The methods for determining the structure of novel POMs is perfectly established as well as the techniques to investigate their stability in solution. For this purpose, the typical techniques are the single-crystal X-ray diffraction for the solid-state characterization and the multinuclear high-resolution NMR spectroscopy for the analysis of the solutions.


Nevertheless, with the ever-increasing complexity of the supramolecular POM-

based systems, the limits of the NMR techniques can be rapidly reached when the size of the aggregates increases and the viscosity of the medium decreases, as in hydrogels and supramolecular polymers. In this case, other techniques more suited to semi-fluids and soft matter systems, such as cryo-TEM, AFM, Small-angle X-ray or neutron scattering (SAXS or SANS) can take over. One of the goals of this axis will be to explore the possibilities and the limits of such techniques.



Energy applications, sustainable and environmental chemistry

The challenges that POM chemistry is facing are therefore manifold. The IRN support efforts to stimulate discussions about new or already known catalytic reactions, possibly combined with



mechanistic investigations based on

Reagent



Product

experimental or computational studies. While electro-assisted water reduction by POMs is well illustrated, utilization of CO₂ for redox functionalization of organic substrates, CO₂ reduction to fuels, biomass conversion and even natural gas transformation (through photochemically assisted C-H bond activation) remain more challenging and require greater efforts. To break down the scientific barriers, it will be necessary to combine expertise in catalysis, photochemistry, electrochemistry, surface science, modelling, chemical engineering and as already stated above, data science.

POM engineering for functional materials

Electro-active molecules that can be switched from one redox state to another have been proposed for the development of molecular memory devices. In order to move from the status of a potential application to the preparation of well-defined models (devices) that are needed to assess the benefits of using POMs, a number of synthetic challenges still need to be overcome. It is therefore essential to

control the POM shape processing on the solid surface (interfacial contact or electrodes) and to use and develop accurate physical techniques to characterize the electrical properties. This will require a high level of engineering of the molecules/electrode interface and therefore the establishment of close collaboration with physicists, industrial partners and also theoreticians to gain some understanding about the mechanisms of charge transfer, transport or even storage. These are some of the main challenges for POM chemists at the frontier with materials chemistry.

